

Quantum Affine Algebra and Universal R -Matrix with Spectral Parameter

Yao-Zhong Zhang and Mark D. Gould

Department of Mathematics, University of Queensland, Brisbane, Qld 4072, Australia

Abstract:

Using the previously obtained universal R -matrix for the quantized nontwisted affine Lie algebras $U_q(A_1^{(1)})$ and $U_q(A_2^{(1)})$, we determine the explicitly spectral-dependent universal R -matrix for the corresponding quantum Lie algebras $U_q(A_1)$ and $U_q(A_2)$. As their applications, we reproduce the well-known results in the fundamental representations and we also derive an extremely explicit formula of the spectral-dependent R -matrix for the adjoint representation of $U_q(A_2)$, the simplest non-trivial case when the tensor product decomposition of the representation with itself has finite multiplicity.

Mathematics Subject Classifications (1991): 81R10, 17B37, 16W30

1. Introduction: In this Letter, we present two main results: (i) the universal R -matrix for quantum Lie algebras $U_q(A_1)$ and $U_q(A_2)$ with explicit spectral parameter dependence; (ii) an extremely explicit formula for the spectral-dependent R -matrix in the adjoint representation of $U_q(A_2)$, the simplest nontrivial case when the tensor product decomposition of the representation with itself has finite multiplicity.

Since their invention, quantum algebras [1][2] [3] have been employed, as a main technique, to find spectral parameter dependent solution to quantum Yang-Baxter equation (QYBE) (that is spectral-dependent R -matrix). Two conventional approaches to this effect in literatures may be viewed as the following two seemingly different but essentially related "Yang-Baxterization" processes: (i) one starts from a quantum simple Lie algebra, affinizes it, thus giving rise to the Jimbo equations, and then solves the Jimbo equations [2][5]; (ii) one starts with some braid group representation of a quantum simple Lie algebra and then tries to introduce a spectral parameter in such a way that the spectral-dependent QYBE is satisfied [4]. The "abelian Yang-Baxterization", where the tensor product decomposition of representation with itself is multiplicity-free, is extensively studied in literatures for lower representations. For higher representations, the "fusion" procedure is suggested which is not, however, easily workable. As to "non-abelian Yang-Baxterization" in which the tensor product decomposition of representation with itself has finite multiplicity, very few attempt has currently been made, due to conceivable complexities caused by the appearance of finite multiplicity. Therefore, it seems desirable to develop more effective procedures to cope with the situations.

In this Letter we will present a new way of obtaining spectral-dependent R -matrix for quantum Lie algebras. Our idea reverses, in some sense, the above process. Namely, we start from the universal R -matrix of quantum affine Lie algebra $U_q(\mathcal{G}^{(1)})$ and then apply it to finite-dimensional loop representations $V(z)$ of $U_q(\mathcal{G}^{(1)})$, which is isomorphic to the ones $V \otimes \mathbf{C}(z, z^{-1})$ of the corresponding quantum simple Lie algebra $U_q(\mathcal{G})$. In this way, a spectral parameter appears automatically and we obtain the spectral-dependent solution to QYBE for the latter. A remarkable advantage lying this approach is that both the case of multiplicity-free and the case with finite multiplicity can be treated in a unified fashion. As a matter of fact, we are able to get a spectral-dependent *universal* form of R -matrix for $U_q(A_1)$ and $U_q(A_2)$, regardless of

representations and multiplicities considered. Applying to some concrete representations, we are able to reproduce the well-known results in the fundamental representation and to obtain an extremely explicit formula for R -matrix of $U_q(A_2)$ for the adjoint representation.

2. Universal R -Matrix for $U_q(A_1^{(1)})$ and $U_q(A_2^{(1)})$: For self-contained, this section is devoted to a brief and quick review of the construction of the universal R -matrix for $U_q(A_1^{(1)})$ [6] and for $U_q(A_2^{(1)})$ [6][7]. Throughout the paper, we use the notations: $(\text{ad}_q x_\alpha)x_\beta = [x_\alpha, x_\beta] = x_\alpha x_\beta - q^{(\alpha, \beta)} x_\beta x_\alpha$, $\theta(q^h) = q^{-h}$, $\theta(E_i) = F_i$, $\theta(F_i) = E_i$, $\theta(q) = q^{-1}$, $(n)_q = (1 - q^n)/(1 - q)$, $[n]_q = (q^n - q^{-n})/(q - q^{-1})$, $q_\alpha = q^{(\alpha, \alpha)}$, $\exp_q(x) = \sum_{n \geq 0} \frac{x^n}{(n)_q!}$, $(n)_q! = (n)_q(n-1)_q \dots (1)_q$.

We start with rank 2 case. Fix the normal ordering in the positive root system Δ_+ of $A_1^{(1)}$: $\alpha, \alpha + \delta, \dots, \alpha + n\delta, \dots, \delta, 2\delta, \dots, m\delta, \dots, (\delta - \alpha) + l\delta, \dots, \delta - \alpha$, where $\alpha, \delta - \alpha$ are simple roots and δ is the minimal positive imaginary root. One has [6] the universal R -matrix for $U_q(A_1^{(1)})$:

$$R = \left(\prod_{n \geq 0} \exp_{q_\alpha}((q - q^{-1})(E_{\alpha+n\delta} \otimes F_{\alpha+n\delta})) \right) \cdot \exp \left(\sum_{n > 0} n [n]_{q_\alpha}^{-1} (q_\alpha - q_\alpha^{-1})(E_{n\delta} \otimes F_{n\delta}) \right) \cdot \left(\prod_{n \geq 0} \exp_{q_\alpha}((q - q^{-1})(E_{(\delta-\alpha)+n\delta} \otimes F_{(\delta-\alpha)+n\delta})) \right) \cdot q^{\frac{1}{2}h_\alpha \otimes h_\alpha + c \otimes d + d \otimes c} \quad (1)$$

where $c = h_\alpha + h_{\delta-\alpha}$ and the Cartan-Weyl generators, $E_\gamma, F_\gamma = \theta(E_\gamma)$, $\gamma \in \Delta_+$, are given by: $\tilde{E}_\delta = [(\alpha, \alpha)]_q^{-1} [E_\alpha, E_{\delta-\alpha}]_q$, $E_{\alpha+n\delta} = (-1)^n (\text{ad} \tilde{E}_\delta)^n E_\alpha$, $E_{(\delta-\alpha)+n\delta} = (\text{ad} \tilde{E}_\delta)^n E_{\delta-\alpha}$, $\tilde{E}_{n\delta} = [(\alpha, \alpha)]_q^{-1} [E_{\alpha+(n-1)\delta}, E_{\delta-\alpha}]_q$ and

$$\tilde{E}_{n\delta} = \sum_{p_1+2p_2+\dots+np_n=n} \frac{(q^{(\alpha, \alpha)} - q^{-(\alpha, \alpha)})^{\sum_i p_i - 1}}{p_1! \dots p_n!} (E_\delta)^{p_1} (E_{2\delta})^{p_2} \dots (E_{n\delta})^{p_n} \quad (2)$$

The order in the product (1) concides with the above chosen normal order.

We now consider rank 3 case. Fix the order in positive root system Δ_+ of $A_2^{(1)}$: $\alpha, \alpha + \delta, \dots, \alpha + m_1\delta, \dots, \alpha + \beta, \alpha + \beta + \delta, \dots, \alpha + \beta + m_2\delta, \dots, \beta, \beta + \delta, \dots, \beta + m_3\delta, \dots, \delta, 2\delta, \dots, k\delta, \dots, (\delta - \beta) + l_1\delta, \dots, \delta - \beta, \dots, (\delta - \alpha) + l_2\delta, \dots, \delta - \alpha, \dots, (\delta - \alpha - \beta) + l_3\delta, \dots, \delta - \alpha - \beta$, where $m_i, k, l_i \geq 0$, $i = 1, 2, 3$. Then one can show [6][7] (see, in particular, [7]) that the universal R -matrix for $U_q(A_2^{(1)})$ takes the explicit form

$$R = \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1})(E_{\alpha+n\delta} \otimes F_{\alpha+n\delta}) \right)$$

$$\begin{aligned}
& \cdot \prod_{n \geq 0} \exp_{q_{\alpha+\beta}} \left((q - q^{-1})(E_{\alpha+\beta+n\delta} \otimes F_{\alpha+\beta+n\delta}) \right) \\
& \cdot \prod_{n \geq 0} \exp_{q_\beta} \left((q - q^{-1})(E_{\beta+n\delta} \otimes F_{\beta+n\delta}) \right) \\
& \cdot \exp \left(\sum_{n > 0} \sum_{i,j=1}^2 C_{ij}^n(q) (q - q^{-1})(E_{n\delta}^{(i)} \otimes F_{n\delta}^{(j)}) \right) \\
& \cdot \prod_{n \geq 0} \exp_{q_\beta} \left((q - q^{-1})(E_{(\delta-\beta)+n\delta} \otimes F_{(\delta-\beta)+n\delta}) \right) \\
& \cdot \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1})(E_{(\delta-\alpha)+n\delta} \otimes F_{(\delta-\alpha)+n\delta}) \right) \\
& \cdot \prod_{n \geq 0} \exp_{q_{\alpha+\beta}} \left((q - q^{-1})(E_{(\delta-\alpha-\beta)+n\delta} \otimes F_{(\delta-\alpha-\beta)+n\delta}) \right) \\
& \cdot q^{\sum_{i,j=1}^2 (a_{\text{sym}}^{-1})^{ij} h_i \otimes h_j + c \otimes d + d \otimes c}
\end{aligned} \tag{3}$$

where $c = h_0 + h_\psi$ with $\psi = \alpha + \beta$ being the highest root of $A_2^{(1)}$ and $(a_{\text{sym}}^{ij}) = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$;

$$(C_{ij}^n(q)) = (C_{ji}^n(q)) = \frac{n}{[n]_q} \frac{[2]_q^2}{q^{2n} + 1 + q^{-2n}} \begin{pmatrix} q^n + q^{-n} & (-1)^n \\ (-1)^n & q^n + q^{-n} \end{pmatrix} \tag{4}$$

and the Cartan-Weyl generators, E_γ , $F_\gamma = \theta(E_\gamma)$, $\gamma \in \Delta_+$, are given by ($\alpha_i = \alpha, \beta, \alpha + \beta$ below) : $E_{\alpha+\beta} = [E_\alpha, E_\beta]_q$, $E_{\delta-\alpha} = [E_\beta, E_{\delta-\alpha-\beta}]_q$, $E_{\delta-\beta} = [E_\alpha, E_{\delta-\alpha-\beta}]_q$, $\tilde{E}_\delta^{(i)} = [(\alpha_i, \alpha_i)]_q^{-1} [E_{\alpha_i}, E_{\delta-\alpha_i}]_q$, $E_{\alpha_i+n\delta} = (-1)^n (\text{ad} \tilde{E}_\delta^{(i)})^n E_{\alpha_i}$, $E_{\delta-\alpha_i+n\delta} = (\text{ad} \tilde{E}_\delta^{(i)})^n E_{\delta-\alpha_i}$, $\tilde{E}_{n\delta}^{(i)} = [(\alpha_i, \alpha_i)]_q^{-1} [E_{\alpha_i+(n-1)\delta}, E_{\delta-\alpha_i}]_q$ and

$$\tilde{E}_{n\delta}^{(i)} = \sum_{p_1+2p_2+\dots+np_n=n} \frac{(q^{(\alpha_i, \alpha_i)} - q^{-(\alpha_i, \alpha_i)})^{\sum_i p_i - 1}}{p_1! \dots p_n!} (E_\delta^{(i)})^{p_1} (E_{2\delta}^{(i)})^{p_2} \dots (E_{n\delta}^{(i)})^{p_n} \tag{5}$$

The order in the product of (3) concides with the above defined order.

3. Universal R -matrix with Spectral Parameter: We come to our main concerns in this Letter. We only list our results and details will be published elsewhere [8].

It can be shown that for any $z \in \mathbf{C}^\times$, there is a homomorphism of algebras $\text{ev}_z: U_q(A_1^{(1)}) \rightarrow U_q(A_1)$ given by: $\text{ev}_z(E_\alpha) = E_\alpha$, $\text{ev}_z(F_\alpha) = F_\alpha$, $\text{ev}_z(h_\alpha) = h_\alpha$, $\text{ev}_z(c) = 0$, $\text{ev}_z(E_{\delta-\alpha}) = zF_\alpha$, $\text{ev}_z(F_{\delta-\alpha}) = z^{-1}E_\alpha$, $\text{ev}_z(h_{\delta-\alpha}) = -h_\alpha$. Then from eq.(1) one obtains the spectral-dependent universal R -matrix, $R(x, y) \equiv (\text{ev}_x \otimes \text{ev}_y)R$, for $U_q(A_1)$:

$$R(x, y) = \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1}) \left(\frac{x}{y} \right)^n \left(q^{-nh_\alpha} E_\alpha \otimes F_\alpha q^{nh_\alpha} \right) \right)$$

$$\begin{aligned}
& \cdot \exp \left(\sum_{n>0} n[n]_{q_\alpha}^{-1} (q_\alpha - q_\alpha^{-1}) \left(\frac{x}{y} \right)^n (E'_{n\delta} \otimes F'_{n\delta}) \right) \\
& \cdot \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1}) \left(\frac{x}{y} \right)^{n+1} (F_\alpha q^{-nh_\alpha} \otimes q^{nh_\alpha} E_\alpha) \right) \cdot q^{\frac{1}{2}h_\alpha \otimes h_\alpha} \quad (6)
\end{aligned}$$

where $E'_{n\delta}$ and $F'_{n\delta}$ are determined by the following equalities of formal power series:

$$\begin{aligned}
1 + (q_\alpha - q_\alpha^{-1}) \sum_{k=1}^{\infty} \tilde{E}'_{k\delta} u^k &= \exp \left((q_\alpha - q_\alpha^{-1}) \sum_{l=1}^{\infty} E'_{l\delta} u^l \right) \\
1 - (q_\alpha - q_\alpha^{-1}) \sum_{k=1}^{\infty} \tilde{F}'_{k\delta} u^{-k} &= \exp \left(-(q_\alpha - q_\alpha^{-1}) \sum_{l=1}^{\infty} F'_{l\delta} u^{-l} \right) \quad (7)
\end{aligned}$$

in which $\tilde{E}'_{n\delta} = [2]_q^{-1} (-1)^{n-1} q^{-(n-1)h_\alpha} (E_\alpha F_\alpha - q^{-2n} F_\alpha E_\alpha)$, $\tilde{F}'_{n\delta} = \theta(\tilde{E}'_{n\delta})$.

We now apply (6) to $V_{1/2} \otimes V_{1/2}$, where $V_{1/2}$ is the fundamental representation of $U_q(A_1)$.

One can show that in this case

$$R_{1/2,1/2}(x,y) = f_q(x,y) \cdot \left(e_{11} + e_{44} + \frac{q^{-1}(y-x)}{y-q^{-2}x} (e_{22} + e_{33}) + \frac{1-q^{-2}}{y-q^{-2}x} (ye_{23} + xe_{32}) \right) \quad (8)$$

where (and below) e_{ij} is the matrix satisfying $(e_{ij})_{kl} = \delta_{ik} \delta_{jl}$ and

$$f_q(x,y) = q^{1/2} \cdot \exp \left(\sum_{n>0} \frac{q^n - q^{-n}}{q^n + q^{-n}} \frac{(x/y)^n}{n} \right) \quad (9)$$

We thus reproduce the well-known result [2], up to the scalar factor $f_q(x,y)$. In [6] KT obtained (8) directly from (1).

We then turn to $U_q(A_2^{(1)})$. In this case, one may show that for any $z \in \mathbf{C}^\times$, there is a homomorphism of algebras $\text{ev}_z: U_q(A_2^{(1)}) \rightarrow U_q(A_2)$ given by: $\text{ev}_z(E_\alpha) = E_\alpha$, $\text{ev}_z(F_\alpha) = F_\alpha$, $\text{ev}_z(h_\alpha) = h_\alpha$, $\text{ev}_z(E_\beta) = E_\beta$, $\text{ev}_z(F_\beta) = F_\beta$, $\text{ev}_z(h_\beta) = h_\beta$, $\text{ev}_z(E_{\delta-\alpha-\beta}) = zF_{\alpha+\beta}q^{(h_\beta-h_\alpha)/3}$, $\text{ev}_z(F_{\delta-\alpha-\beta}) = z^{-1}q^{(h_\alpha-h_\beta)/3}E_{\alpha+\beta}$, $\text{ev}_z(h_{\delta-\alpha-\beta}) = -h_{\alpha+\beta}$, $\text{ev}_z(c) = 0$. Thus, from eq.(3) one is able to deduce the explicitly spectral-dependent universal R -matrix for $U_q(A_2)$, $R(x,y) \equiv (\text{ev}_x \otimes \text{ev}_y)R$, given by

$$\begin{aligned}
R(x,y) &= \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1}) \left(\frac{x}{y} \right)^n (q^{-nh_\alpha} E_\alpha q^{-n(h_\alpha+2h_\beta)/3} \otimes q^{n(h_\alpha+2\beta)/3} F_\alpha q^{nh_\alpha}) \right) \\
&\cdot \prod_{n \geq 0} \exp_{q_{\alpha+\beta}} \left((q - q^{-1}) \left(\frac{x}{y} \right)^n (q^{-nh_{\alpha+\beta}} E_{\alpha+\beta} q^{n(h_\beta-h_\alpha)/3} \otimes q^{n(h_\alpha-\beta)/3} F_{\alpha+\beta} q^{nh_{\alpha+\beta}}) \right) \\
&\cdot \prod_{n \geq 0} \exp_{q_\beta} \left((q - q^{-1}) \left(\frac{x}{y} \right)^n (E'_{\beta+n\delta} \otimes F'_{\beta+n\delta}) \right) \\
&\cdot \exp \left(\sum_{n>0} \sum_{i,j=1}^2 C_{ij}^n(q) (q - q^{-1}) \left(\frac{x}{y} \right)^n (E'_{n\delta}^{(i)} \otimes F'_{n\delta}^{(j)}) \right)
\end{aligned}$$

$$\begin{aligned}
& \cdot \prod_{n \geq 0} \exp_{q_\beta} \left((q - q^{-1}) \left(\frac{x}{y} \right)^{n+1} \left(E'_{(\delta-\beta)+n\delta} \otimes F'_{(\delta-\beta)+n\delta} \right) \right) \\
& \cdot \prod_{n \geq 0} \exp_{q_\alpha} \left((q - q^{-1}) \left(\frac{x}{y} \right)^{n+1} \left(q^{-(n+1)(h_\alpha+2h_\beta)/3} F_\alpha q^{-nh_\alpha} \otimes q^{nh_\alpha} E_\alpha q^{(n+1)(h_\alpha+2h_\beta)/3} \right) \right) \\
& \cdot \prod_{n \geq 0} \exp_{q_{\alpha+\beta}} \left((q - q^{-1}) \left(\frac{x}{y} \right)^{n+1} \left(q^{(n+1)(h_\beta-h_\alpha)/3} F_{\alpha+\beta} q^{-nh_{\alpha+\beta}} \right. \right. \\
& \quad \left. \left. \otimes q^{nh_{\alpha+\beta}} E_{\alpha+\beta} q^{(n+1)(h_\alpha-h_\beta)/3} \right) \right) \cdot q^{\sum_{i,j=1}^2 (a_{\text{sym}}^{-1})^{ij} h_i \otimes h_j} \quad (10)
\end{aligned}$$

where

$$\begin{aligned}
E'_{\beta+n\delta} &= (-1)^n [2]_q^{-n} q^n \left\{ \left(\text{ad}'_{q^{-1}} \mathcal{E} \right)^n E_\beta \right\} q^{n(h_\beta-h_\alpha)/3} \\
F'_{\beta+n\delta} &= [2]_q^{-n} q^{n(h_\alpha-h_\beta)/3} \left(\text{ad}'_{q^{-1}} \mathcal{F} \right)^n F_\beta \\
E'_{(\delta-\beta)+n\delta} &= [2]_q^{-n} q^{-n} \left\{ \left(\text{ad}'_q \mathcal{E} \right)^n \left(\text{ad}'_{q^{-2}} E_\alpha \right) F_{\alpha+\beta} \right\} q^{(n+1)(h_\beta-h_\alpha)/3} \\
F'_{(\delta-\beta)+n\delta} &= (-1)^n [2]_q^{-n} q^{(n+1)(h_\alpha-h_\beta)/3} \left(\text{ad}'_q \mathcal{F} \right)^n \left(\text{ad}'_{q^2} E_{\alpha+\beta} \right) F_\alpha \quad (11)
\end{aligned}$$

$\mathcal{E} = (\text{ad}'_{q^{-1}} E_\beta)(\text{ad}'_{q^{-2}} E_\alpha) F_{\alpha+\beta}$, $\mathcal{F} = (\text{ad}'_q (\text{ad}'_{q^2} E_{\alpha+\beta}) F_\alpha) F_\beta$ and $(\text{ad}'_{Q\mathcal{A}}) \cdot \mathcal{B} \equiv \mathcal{AB} - Q\mathcal{BA}$; $E'_{n\delta}^{(i)}$ and $F'_{n\delta}^{(i)}$ are determined by the equalities of formal series: $(\alpha_i = \alpha, \beta)$

$$\begin{aligned}
1 + (q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{k=1}^{\infty} \tilde{E}_{k\delta}^{(i)} u^k &= \exp \left((q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{l=1}^{\infty} E_{l\delta}^{(i)} u^l \right) \\
1 - (q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{k=1}^{\infty} \tilde{F}_{k\delta}^{(i)} u^{-k} &= \exp \left(-(q_{\alpha_i} - q_{\alpha_i}^{-1}) \sum_{l=1}^{\infty} F_{l\delta}^{(i)} u^{-l} \right) \quad (12)
\end{aligned}$$

in which

$$\begin{aligned}
\tilde{E}_{n\delta}^{(\alpha)} &= (-1)^{n-1} [2]_q^{-1} (E_\alpha F_\alpha - q^{-2n} F_\alpha E_\alpha) q^{-(n-1)h_\alpha} q^{-n(h_\alpha+2h_\beta)/3} \\
\tilde{F}_{n\delta}^{(\alpha)} &= (-1)^{n-1} [2]_q^{-1} q^{(n-1)h_\alpha} q^{n(h_\alpha+2h_\beta)/3} (F_\alpha E_\alpha - q^{2n} E_\alpha F_\alpha) \\
\tilde{E}_{n\delta}^{(\beta)} &= (-1)^n [2]_q^{-n} q^{n-2} \left\{ \left(\text{ad}'_{q^{-n+2}} \mathcal{F}' \right) \cdot \left(\text{ad}'_{q^{-1}} \mathcal{E} \right)^{n-1} E_\beta \right\} q^{n(h_\beta-h_\alpha)/3} \\
\tilde{F}_{n\delta}^{(\beta)} &= [2]_q^{-n} q^{n-1} q^{n(h_\alpha-h_\beta)/3} \left(\text{ad}'_{q^{-n+2}} \mathcal{E}' \right) \cdot \left(\text{ad}'_{q^{-1}} \mathcal{F} \right)^{n-1} F_\beta \\
\mathcal{E}' &= E_{\alpha+\beta} F_\alpha - q^2 F_\alpha E_{\alpha+\beta}, \quad \mathcal{F}' = E_\alpha F_{\alpha+\beta} - q^{-2} F_{\alpha+\beta} E_\alpha \quad (13)
\end{aligned}$$

We apply (10) to $V_{(3)} \otimes V_{(3)}$ in which $V_{(3)}$ denotes the fundamental representation of $U_q(A_2)$.

One can show that (10) gives rise to

$$\begin{aligned}
R_{(3),(3)}(x, y) &= f_q(x, y) \cdot \left(e_{11} + e_{55} + e_{99} + \frac{q^{-1}(y-x)}{y-q^{-2}x} (e_{22} + e_{33} + e_{44} + e_{66} + e_{77} + e_{88}) + \right. \\
&\quad \left. + \frac{y(1-q^{-2})}{y-q^{-2}x} (e_{24} + e_{37} + e_{68}) + \frac{x(1-q^{-2})}{y-q^{-2}x} (e_{42} + e_{73} + e_{86}) \right) \quad (14)
\end{aligned}$$

where

$$f_q(x, y) = q^{2/3} \cdot \exp \left(\sum_{n>0} \frac{q^{2n} - q^{-2n}}{q^{2n} + 1 + q^{-2n}} \frac{(x/y)^n}{n} \right) \quad (15)$$

Eq.(14) is nothing but the well-known result [2], up to the scalar factor $f_q(x, y)$.

We then apply (10) to a very interesting case: $V_{(8)} \otimes V_{(8)}$, where $V_{(8)}$ stands for the adjoint representation of $U_q(A_2)$. This is simplest nontrivial example where the tensor product decomposition is with finite multiplicity. It can be shown that spectral-dependent R -matrix for the adjoint representation takes the following compact and explicit form,

$$\begin{aligned} R_{(8),(8)}(x, y) = & \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^n (E'_{\alpha+n\delta} \otimes F'_{\alpha+n\delta}) + y^2 f(e_{36} \otimes e_{63}) \right\} \\ & \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^n (E'_{\alpha+\beta+n\delta} \otimes F'_{\alpha+\beta+n\delta}) + y^2 f(e_{18} \otimes e_{81}) \right\} \\ & \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^n (E'_{\beta+n\delta} \otimes F'_{\beta+n\delta}) + y^2 f'(e_{27} \otimes e_{72}) \right\} \\ & \cdot \{ \text{imaginary root vectors contribution} \} \\ & \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^{n+1} (E'_{(\delta-\beta)+n\delta} \otimes F'_{(\delta-\beta)+n\delta}) + x^2 g'(e_{72} \otimes e_{27}) \right\} \\ & \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^{n+1} (E'_{(\delta-\alpha)+n\delta} \otimes F'_{(\delta-\alpha)+n\delta}) + x^2 g(e_{63} \otimes e_{36}) \right\} \\ & \cdot \left\{ 1 + (q - q^{-1}) \sum_{n=0}^{\infty} \left(\frac{x}{y} \right)^{n+1} (E'_{(\delta-\alpha-\beta)+n\delta} \otimes F'_{(\delta-\alpha-\beta)+n\delta}) + x^2 g(e_{81} \otimes e_{18}) \right\} \\ & \cdot \left\{ q^2 \sum_{i=1}^8 (1 - \delta_{i4} - \delta_{i5})(e_{ii} \otimes e_{ii}) + q(e_{11} \otimes e_{22} + e_{11} \otimes e_{33} + e_{22} \otimes e_{66} + \right. \\ & + e_{33} \otimes e_{77} + e_{66} \otimes e_{88} + e_{77} \otimes e_{88} + \{\leftrightarrow\}) + q^{-1}(e_{11} \otimes e_{66} + e_{11} \otimes e_{77} + \\ & + e_{22} \otimes e_{33} + e_{22} \otimes e_{88} + e_{33} \otimes e_{88} + e_{66} \otimes e_{77} + \{\leftrightarrow\}) + \\ & \left. + q^{-2}(e_{11} \otimes e_{88} + e_{22} \otimes e_{77} + e_{33} \otimes e_{66} + \{\leftrightarrow\}) \right\} \end{aligned} \quad (16)$$

where $f = [2]_q q^{-1}(q - q^{-1})^2 (y + q^4 x) / \{(y^2 - x^2)(y - q^2 x)\}$, $g = [2]_q q^{-1}(q - q^{-1})^2 (y + q^{-2} x) / \{(y^2 - x^2)(y - q^{-2} x)\}$, $f' = [2]_q^{-2}(q - q^{-1})^2 (y^2 - x^2)^{-1} (q^{-1}[2]_q^3 + F(q; x, y))$, $g' = [2]_q^{-2}(q - q^{-1})^2 (y^2 - x^2)^{-1} (q^{-1}[2]_q^3 + F(q^{-1}; x, y))$, $F(q, x, y) = q^{-4}x/(y - q^{-4}x) + 2[3]_q x/(y - x) + [3]_q^2 q^4 x/(y - q^4 x)$, and " $\{\leftrightarrow\}$ " denotes the interchange to $X \otimes Y$ of the quantities within the same bracket;

$$\begin{aligned} E'_{\alpha+n\delta} &= (-1)^n \left\{ q^{-2n} e_{12} + [2]_q^{1/2} q^{-2n} e_{34} + [2]_q^{1/2} e_{46} + e_{78} \right\} \\ F'_{\alpha+n\delta} &= (-1)^n \left\{ q^{2n} e_{21} + [2]_q^{1/2} q^{2n} e_{43} + [2]_q^{1/2} e_{64} + e_{87} \right\} \\ E'_{\alpha+\beta+n\delta} &= (-1)^n \left\{ -[2]_q^{-1/2} q^{-2n-2} e_{14} + ([3]_q/[2]_q)^{1/2} q^{-2n} e_{15} - q^{-1} e_{26} + \right. \end{aligned}$$

$$\begin{aligned}
& + q^{-2n} e_{37} + [2]_q^{-1/2} q e_{48} - ([3]_q/[2]_q)^{1/2} q^{-1} e_{58} \Big\} \\
F'_{\alpha+\beta+n\delta} &= (-1)^n \Big\{ -[2]_q^{-1/2} q^{2n+2} e_{41} + ([3]_q/[2]_q)^{1/2} q^{2n} e_{51} - q e_{62} + \\
& + q^{2n} e_{73} + [2]_q^{-1/2} q^{-1} e_{84} - ([3]_q/[2]_q)^{1/2} q e_{85} \Big\} \\
E'_{\beta+n\delta} &= q^n e_{13} + [2]_q^{-1/2} q^n e_{24} + ([3]_q/[2]_q)^{1/2} q^{-3n} e_{25} + \\
& + [2]_q^{-1/2} q^{-3n} e_{47} + ([3]_q/[2]_q)^{1/2} q^n e_{57} + q^{-3n} e_{68} \\
F'_{\beta+n\delta} &= q^{-n} e_{31} + [2]_q^{-1/2} q^{-n} e_{42} + ([3]_q/[2]_q)^{1/2} q^{3n} e_{52} + \\
& + [2]_q^{-1/2} q^{3n} e_{74} + ([3]_q/[2]_q)^{1/2} q^{-n} e_{75} + q^{3n} e_{86} \\
E'_{(\delta-\beta)+n\delta} &= -q^{n+2} e_{31} - [2]_q^{-1/2} q^{n+3} e_{42} - ([3]_q/[2]_q)^{1/2} q^{-3n-1} e_{52} \\
& - [2]_q^{-1/2} q^{-3n-3} e_{74} - ([3]_q/[2]_q)^{1/2} q^{n+1} e_{75} - q^{-3n-2} e_{86} \\
F'_{(\delta-\beta)+n\delta} &= -q^{-n-2} e_{13} - [2]_q^{-1/2} q^{-n-3} e_{24} - ([3]_q/[2]_q)^{1/2} q^{3n+1} e_{25} \\
& - [2]_q^{-1/2} q^{3n+3} e_{47} - ([3]_q/[2]_q)^{1/2} q^{-n-1} e_{57} - q^{3n+2} e_{68} \\
E'_{(\delta-\alpha)+n\delta} &= (-1)^n \Big\{ q^{-2n-1} e_{21} + [2]_q^{1/2} q^{-2n} e_{43} + [2]_q^{1/2} e_{64} + q e_{87} \Big\} \\
F'_{(\delta-\alpha)+n\delta} &= (-1)^n \Big\{ q^{2n+1} e_{12} + [2]_q^{1/2} q^{2n} e_{34} + [2]_q^{1/2} e_{46} + q^{-1} e_{78} \Big\} \\
E'_{(\delta-\alpha-\beta)+n\delta} &= (-1)^n \Big\{ -[2]_q^{-1/2} q^{-2n+2} e_{41} + ([3]_q/[2]_q)^{1/2} q^{-2n} e_{51} - q^2 e_{62} \\
& + q^{-2n-1} e_{73} + [2]_q^{-1/2} q^{-1} e_{84} - ([3]_q/[2]_q)^{1/2} q e_{85} \Big\} \\
F'_{(\delta-\alpha-\beta)+n\delta} &= (-1)^n \Big\{ -[2]_q^{-1/2} q^{2n-2} e_{14} + ([3]_q/[2]_q)^{1/2} q^{2n} e_{15} - q^{-2} e_{26} + \\
& + q^{2n+1} e_{37} + [2]_q^{-1/2} q e_{48} - ([3]_q/[2]_q)^{1/2} q^{-1} e_{58} \Big\} \\
\{\text{imaginary root vectors contribution}\} &= a'/a \sum_{i=1}^8 (1 + (b/b' - 1)\delta_{i4} + (c/c' - 1)\delta_{i5}) (e_{ii} \otimes e_{ii}) + \\
& + a'(e_{11} \otimes e_{22} + e_{11} \otimes e_{33}) + aa'(e_{11} \otimes e_{44}) + a'c(e_{11} \otimes e_{55}) + a(e_{11} \otimes e_{66}) + \\
& + aa'c(e_{11} \otimes e_{77}) + ac(e_{11} \otimes e_{88}) + 1/a(e_{22} \otimes e_{11}) + 1/b'(e_{22} \otimes e_{33}) + \\
& + a'/b'(e_{22} \otimes e_{44}) + a'c(e_{22} \otimes e_{55}) + a'(e_{22} \otimes e_{66}) + aa'c/b'(e_{22} \otimes e_{77}) + \\
& + aa'c(e_{22} \otimes e_{88}) + 1/a(e_{33} \otimes e_{11}) + b(e_{33} \otimes e_{22}) + a'b(e_{33} \otimes e_{44}) + \\
& + ab(e_{33} \otimes e_{66}) + a'(e_{33} \otimes e_{77}) + a(e_{33} \otimes e_{88}) + 1/(aa')(e_{44} \otimes e_{11}) + \\
& + b/a(e_{44} \otimes e_{22}) + 1/(ab')(e_{44} \otimes e_{33}) + a'b(e_{44} \otimes e_{66}) + a'/b'(e_{44} \otimes e_{77}) + \\
& + aa'(e_{44} \otimes e_{88}) + 1/(ac')(e_{55} \otimes e_{11} + e_{55} \otimes e_{22}) + a'c(e_{55} \otimes e_{77} + e_{55} \otimes e_{88}) + \\
& + 1/a'(e_{66} \otimes e_{11}) + 1/a(e_{66} \otimes e_{22}) + 1/(a'b')(e_{66} \otimes e_{33}) + 1/(ab')(e_{66} \otimes e_{44}) +
\end{aligned}$$

$$\begin{aligned}
& + 1/b'(e_{66} \otimes e_{77}) + a'(e_{66} \otimes e_{88}) + 1/(aa'c')(e_{77} \otimes e_{11}) + b/(aa'c')(e_{77} \otimes e_{22}) + \\
& + 1/a(e_{77} \otimes e_{33}) + b/a(e_{77} \otimes e_{44}) + 1/(ac')(e_{77} \otimes e_{55}) + b(e_{77} \otimes e_{66}) + \\
& + a'(e_{77} \otimes e_{88}) + 1/(a'c')(e_{88} \otimes e_{11}) + 1/(aa'c')(e_{88} \otimes e_{22}) + 1/a'(e_{88} \otimes e_{33}) + \\
& + 1/(aa')(e_{88} \otimes e_{44}) + 1/(ac')(e_{88} \otimes e_{55}) + 1/a(e_{88} \otimes e_{66} + e_{88} \otimes e_{77}) \tag{17}
\end{aligned}$$

in which we have defined $a = (y - q^2x)/(y - x)$, $a' = (y - q^{-2}x)/(y - x)$, $b = (y - q^4x)/(y - q^2x)$, $b' = (y - q^{-4}x)/(y - q^{-2}x)$, $c = (y - q^6x)/(y - q^4x)$, $c' = (y - q^{-6}x)/(y - q^{-4}x)$. We remark that (16) is an extremely explicit formula: the sums in (16) can be easily worked out. For compactness, we leave them as the present form.

4. Concluding Remarks: In this letter, we have obtained the spectral-dependent universal R -matrix for $U_q(A_1)$ and $U_q(A_2)$. As their applications, we reproduce from them some well-known results and obtain an extremely explicit and compact formula for spectral-dependent R -matrix in the adjoint representation of $U_q(A_2)$. We believe that our results will be useful in: (i) one dimensional open spin chains [9], (ii) quantization of the conformal affine Liouville (and Toda) theories [10] and (iii) q -deformed WZNW CFT's [11].

One of the authors (Y.Z.Z.) is grateful to Anthony John Bracken for contineous interest, suggestions and discussions, to V.N.Tolstoy for communication of his papers on quantum groups and to R.Cuerno for email correspondences. After completing this paper, Zhong-Qi Ma informed Y.Z.Z. of an earlier paper by him and his coworkers [12] in which part of the results in the present paper – the quantum R -matrix associated with the adjoint reps of $U_q(A_2)$ – was also obtained using the conventional method in the literature – directly solving Jimbo-type equations with the help of quantum CG coefficients (projection operators) of $U_q(A_2)$. The present paper, however, presents a new and different procedure which avoids employing the knowledge of the quantum CG coefficients which are not generally available. Y.Z.Z. thanks Zhong-Qi Ma for e-mail correspondences and for relevant comments. The financial support from Australian Research Council is gratefully acknowledged.

References

- [1] V.G.Drinfeld, *Proc. ICM, Berkeley* **1** (1986) 798
- [2] M.Jimbo, *Lett.Math.Phys.* **10** (1985) 63, *ibid* **11** (1986) 247; *Commun.Math.Phys.* **102** (1986) 247
- [3] N.Reshetikhin, *Quantized universal enveloping algebras, the Yang-Baxter equation and invariants of links: I, II*, preprints LOMI E-4-87, E-17-87
- [4] V.F.R.Jones, *Int.J.Mod.Phys.* **B4** (1990) 701
- [5] R.B.Zhang, M.D.Gould and A.J.Bracken, *Nucl.Phys.* **354** (1991) 625
- [6] S.M.Khoroshkin and U.N.Tolstoy, *Lett.Math.Phys.* **24** (1992) 231; *Funkz.Analiz. i ego Pril.* **26** (1992) 85
- [7] Y.-Z.Zhang and M.D.Gould, *On universal R-Matrix for quantized nontwisted rank 3 affine Lie algebras*, preprint hep-th/9303095, *Lett.Math.Phys.* (in press)
- [8] Y.-Z.Zhang and M.D.Gould, *Quantum affine algebras and universal R-Matrix with spectral parameter, II*, preprint UQMATH-93-06, July 1993, hep-th/9307008
- [9] B.Davies, O.Foda, M.Jimbo, T.Miwa and A.Nakayashiki, *Commun.Math.Phys.* **151** (1993) 89; D.Altschuler and B.Davies, *Quantum loop modules and quantum spin chains*, preprint hep-th/9305135
- [10] O.Babelon and L.Bonora, *Phys.Lett.* **B244** (1990) 220; F.Toppan and Y.-Z.Zhang, *Phys.Lett.* **B292** (1992) 67
- [11] I.B.Frenkel and N.Reshetikhin, *Commun.Math.Phys.* **146** (1992) 1 **180** (1989) 247
- [12] B.-Y.Hou, B.-Y.Hou, Z.-Q.Ma and Y.-D.Yin, *J.Math.Phys.* **32** (1991) 2210